



45A Discovery Way, Acton MA 01720 | Phone: 978-429-2000 | Fax: 978-264-1935 | middlesexdigestive.com

For GLP1 Agonist Medications:

1. Ozempic, Wegovy, Rybelsus (semaglutide)
2. Trulicity (dulaglutide)
3. Mounjaro, Zepbound (tirzepatide)
4. Victoza, Saxenda (liraglutide)
5. Byetta, Bydureon (exanatide)
6. Adlyxin (lixisenatide)

For Endoscopy: The patient can continue their GLP1 medication but they need to start a Full Liquid Diet (no solid foods) 24 hours prior to their procedure time then at midnight prior to their procedure they may have clear liquids only up until 4 hours prior to their endoscopy.

FULL LIQUID DIET- juices(no pulp), milk, ice cream(no solid pieces), popsicles, strained cream soups, Jello, milkshakes, tea, coffee, water, electrolyte drinks or broth. No blending solid foods.

For Colonoscopy: The patient can continue their GLP1 medication - must start clear liquid diet at least 24 hours prior to procedure time. They should already be compliant with this if they follow the colonoscopy instructions. Stop clear liquids 4 hours prior to procedure time.

For SGLT2i Medication

1. Jardiance (empagliflozin)
2. Farxiga (Dapagliflozin)
3. Invokana (canagliflozin)
4. Steglatro (ertugliflozin)
5. Inpefa (sotagliflozin)
6. Steglatro (ertugliflozin)**

For Endoscopy and Colonoscopy: Hold SGLT2i medication for a FULL 3 days (72 hours) prior to their procedure time. The patient can take their last dose 4 days prior to their procedure date. **The exception to this is for those patients on Steglatro (ertugliflozin)-this medication must be held for 4 days/96 hours prior to their procedure time.

If the patient takes more than 2 of these medications for diabetes please have the patient contact their Endocrinologist or PCP as these will need to be held prior to their procedure